


Essential Question

Why is a full shopping cart harder to speed up than an empty one?

Force Causes Acceleration


- Net force causes acceleration
- a points the same way as F_{net}
- No net force, no acceleration



acceleration points with the net force

Force and Acceleration


- More force, more acceleration
- Double F_{net} , double a
- Mass stays the same



straight line through the origin: direct

Mass and Acceleration

- More mass, less acceleration
- Double m , half the a
- Force stays the same



curve that drops: inverse

Newton's Second Law

$$a = F_{\text{net}} / m$$

a

acceleration (m/s²)

F_{net}


net force (N)

m

mass (kg)

Solving for Any Variable


- Find a : $a = F_{\text{net}}/m$
- Find force: $F_{\text{net}} = ma$
- Find mass: $m = F_{\text{net}}/a$



cover the one you want

The Newton


- $1 \text{ N} = 1 \text{ kg} \cdot \text{m/s}^2$
- Gives 1 kg an a of 1 m/s^2
- Units check your work



1 N on 1 kg gives 1 m/s²


Example: Finding Acceleration

A net force of 36 N east acts on a 12 kg wagon. What is the wagon's acceleration?




Example: Finding Net Force

A 75.0 kg sprinter accelerates out of the starting blocks at 4.20 m/s^2 forward. What net force acts on her?




Example: Finding Mass

A net force of 24 N gives a bowling ball an acceleration of 4.0 m/s^2 . What is the ball's mass?



Example: Using a Free-Body Diagram

A 15 kg box is pushed east with a 92 N force. Friction acts on it with 47 N west. What is the box's acceleration?



Big Ideas

- A net force causes acceleration in the same direction
- $a = F_{\text{net}}/m$: more force, more a ; more mass, less a
- Rearrange: $F_{\text{net}} = ma$, $m = F_{\text{net}}/a$
- $1 \text{ N} = 1 \text{ kg} \cdot \text{m}/\text{s}^2$