


Essential Question


How much energy does a moving object have?

Kinetic Energy

- Energy of motion
- Anything moving has KE
- Measured in joules (J)



parked: $KE = 0$



moving: $KE > 0$

The Kinetic Energy Equation

$$KE = \frac{1}{2}mv^2$$

KE

kinetic energy (J)

m

mass (kg)


v

speed (m/s)

$$1 \text{ J} = 1 \text{ kg} \cdot \text{m}^2/\text{s}^2$$


Speed Matters Most

- Double the mass: $2\times$ the KE
- Double the speed: $4\times$ the KE
- Triple the speed: $9\times$ the KE



The Work-Energy Theorem

- Net work changes KE
- $W_{\text{net}} = KE_f - KE_i$
- Positive work speeds it up



$$W_{\text{net}} = \Delta KE$$

Solving for Any Variable

- Write the equation first
- Undo steps in reverse order
- Square root last for v

$$KE = \frac{1}{2}mv^2$$

↓ $\times 2, \div m$

$$v^2 = 2KE / m$$


↓ square root

$$v = \sqrt{(2KE / m)}$$

also: $m = 2KE / v^2$


Example: A Moving Car

A 1,500 kg car drives down the highway at 20.0 m/s. What is its kinetic energy?




Example: Doubling the Speed

A 0.145 kg baseball is thrown at 20.0 m/s, then at 40.0 m/s. Find its kinetic energy at each speed. How do they compare?




Example: Finding Speed

A 60.0 kg skater glides across the ice with 1,920 J of kinetic energy. How fast is she moving?



Example: Work-Energy Theorem

A 2.0 kg cart rolls at 3.0 m/s. A push does 16 J of net work on it in the direction of motion. What is the cart's new speed?



Big Ideas

- $KE = \frac{1}{2}mv^2$: the energy of motion
- Double the speed: 4× the kinetic energy
- Work-energy theorem: $W_{\text{net}} = \Delta KE$
- Find speed: $v = \sqrt{2KE/m}$