

Essential Question

How do Q and K explain Le Châtelier's principle?

A Stress Changes Q or K


Stress	Changes Q?	Changes K?
Add or remove a substance	yes	no
Change volume (gas moles differ)	yes	no
Change temperature	no*	yes

**Q stays the same at first, but K moves away from it.*

Adding a Reactant

- Reactant goes up $\rightarrow$ Q goes down
- $Q < K \rightarrow$ shift right


$$Q = \frac{[\text{products}]}{[\text{reactants}] \uparrow}$$



Adding a Product


- Product goes up $\rightarrow$ Q goes up
- $Q > K \rightarrow$ shift left

$$Q = \frac{[\text{products}] \uparrow}{[\text{reactants}]}$$



Changing Volume and Q

- Halve the volume → every concentration doubles
- The side with more gas moles is affected more



$$Q = \frac{[\text{NO}_2]^2}{[\text{N}_2\text{O}_4]}$$

$$Q_{\text{new}} = \frac{(2[\text{NO}_2])^2}{2[\text{N}_2\text{O}_4]}$$

$$= (4/2) \times Q = 2Q$$



Temperature Changes K

Reaction type	Heat it	Cool it
Endothermic ($\Delta H > 0$)	K increases	K decreases
Exothermic ($\Delta H < 0$)	K decreases	K increases

- Then the system shifts until $Q = K$


Graphing a Stress

- Spike = substance added
- Others adjust gradually




Example: Adding a Reactant

At equilibrium, $[\text{H}_2] = 0.100 \text{ M}$, $[\text{I}_2] = 0.100 \text{ M}$, and $[\text{HI}] = 0.707 \text{ M}$ ($K_c = 50.0$). Extra H_2 is added so $[\text{H}_2]$ instantly becomes 0.200 M . Calculate Q and predict the shift.




Example: Compressing a Gas Mixture

At equilibrium, $[\text{N}_2\text{O}_4] = 0.200 \text{ M}$ and $[\text{NO}_2] = 0.0400 \text{ M}$. The volume is suddenly cut in half. Calculate Q right after and predict the shift.



Example: Heating an Exothermic Reaction

For this reaction, $\Delta H = -57.2 \text{ kJ/mol}$. The temperature is increased. Does K change? Does Q change right away? Which way does the system shift?



Big Ideas

- A stress makes $Q \neq K$
- Concentration or volume: Q changes
- Temperature: K changes
- System shifts until $Q = K$ again

