Name _____ Date _____ Period _____

ZornScience
zornscience.org

7.10 Reaction Quotient and Le Châtelier's Principle

AP Chemistry

A stress makes Q ≠ K. Concentration and volume changes move Q; temperature moves K. The system shifts until Q = K again.

1. For $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$, $K_c = 50.0$. At equilibrium $[H_2] = [I_2] = 0.100$ M and $[HI] = 0.707$ M. I_2 is added so $[I_2] = 0.300$ M. Calculate Q and predict the shift.

2. For $N_2O_4(g) \rightleftharpoons 2NO_2(g)$, $K_c = 0.0200$ with $[N_2O_4] = 0.0800$ M and $[NO_2] = 0.0400$ M. The volume is doubled. Calculate Q right after and predict the shift.

3. For $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$, $\Delta H < 0$. The mixture is heated. What happens to K? To Q right away? Which way does it shift?

4. Argon is added to a mixture of N_2O_4 and NO_2 at constant volume. Does Q change? Does the system shift?

CC BY-NC-SA 4.0 | zornscience.org

5. For $N_2O_4(g) \rightleftharpoons 2NO_2(g)$, $\Delta H = +57.2\ kJ/mol_{rxn}$, complete the table.

Stress	**Q changes?**	**K changes?**	**Shift**
add NO_2			
remove N_2O_4			
decrease the volume			
increase the temperature			

6. Use Q and K to explain why removing a product makes a reaction shift right.

CC BY-NC-SA 4.0 | zornscience.org